



**A STUDY TO ASSESS THE EFFECTIVENESS OF SELF-INSTRUCTIONAL
MODULE ON KNOWLEDGE REGARDING ADVERSE EFFECT OF
THROMBOLYTIC AGENTS ON PATIENT, AMONG STAFF NURSES
WORKING IN CARDIAC UNIT AT SELECTED HOSPITAL OF INDORE, M.P.**

¹Dr. Peter Jasper Youtham, ²Mr. Raj Kumar Gill

¹Research Guide, ²Ph.D. Scholar

Background and purpose:

The time of onset of chest pain was considered as the time of onset of the acute myocardial infarction. Acute myocardial infarction was diagnosed using the standard 12 lead electrocardiogram and cardiac enzyme (World Health Organization criteria). The time from onset of myocardial infarction to treatment was separated into early (<12 h) or late (>12 h); this was the time from initial onset of pain to the time when the patient received treatment. Rapid restoration of patency of the infarct-related artery is the key to preserving myocardium and improving survival in acute myocardial infarction, increase the quality of life of patients, and decrease health expenditure in many countries

In-hospital stroke is associated with slower access to thrombolysis than community-occurring stroke. It has been suggested that lack of knowledge regarding appropriate stroke response among hospital staff may contribute to delays in referral, assessment, and treatment of in-hospital stroke.

Method: A survey was conducted among hospital ward staff members using the adverse effect of thrombolytic agent Questionnaire, which was adapted for use among hospital staff to assess their knowledge on effect of thrombolytic agent, acute treatments, and hospital protocols for treatment of effect of thrombolytic agent.

Results: Ninety-six staff members were interviewed, 81% of whom were clinical staff (medical, nursing, allied health professionals). Ninety-two percent of staff could name ≥ 3 stroke symptoms. Only 49% of staff were aware of thrombolysis treatment, and only 48% could identify the time window for thrombolysis administration, with staff from stroke-related specialties likely to name thrombolysis as an acute treatment for stroke (71%; odds ratio =3.36, 95% confidence interval 1.17-9.61) and identify the correct treatment window (71%; odds ratio =3.55, 95% confidence interval 1.24-10.16). Only 52% of staff on general wards were aware of an in-hospital stroke protocol.

Conclusions: Hospital staff had adequate knowledge on adverse effect of thrombolytic agent; however, there was low awareness of thrombolysis therapy and its correct treatment time window among hospital staff.

References

1. D'Souza, M. S. (2015). Effect of Planned Teaching on knowledge and practice regarding the care of patients receiving Anti-Coagulant Therapy among the Nurses in the Intensive Care Unit in a Selected Hospital, Asian J. Nur. Edu. and Research 5(3): July- Sept.2015; Page 351-362.
2. Mariani, B., Ross, J. G., Paparella, S., & Allen, L. R. (2017). Medication safety simulation to assess student knowledge and competence. Clinical Simulation in Nursing, 13(5), 210-216.
3. Ltheeth, H., & Abbas, S. (2017). Effectiveness of an Educational Program on Nurses' Knowledge Concerning Medication Error at Teaching Hospital in AL-Nasiriyah City. International Journal of Science and Research (IJSR), 6(1).
4. Charalampopoulos, D., Karlis, G., Barouxis, D., Syggelou, A., Mikalli, C., Kountouris, D., & Iacovidou, N. (2016). Theoretical knowledge and skill retention 4 months after a European Paediatric Life Support course. European Journal of Emergency Medicine, 23(1), 56-60.
5. Sneek, S., Isola, A., Saarnio, R. and Boigu, R. (2015). Medication competency of nurses, according to theoretical and drug calculation online exams: A descriptive correlational study. Nurse Education Today, 36 (2016): 195-201.

6. Gorgich, E. A. C., Barfroshan, S., Ghoreishi, G., & Yaghoobi, M. (2016). Investigating the causes of medication errors and strategies to prevention of them from nurses and nursing student viewpoint. *Global journal of health science*, 8(8), 220.
7. Sambu, B. M., Yousif, M. E., & El Mustafa, S.M. (2018). Effect of a Training Program on Nurse's, knowledge, Attitudes and Practice regarding Nursing Care of Acute Myocardial Infarction in Medani Heart Center and Wad Medani Emergency Hospital, Gezira State, Sudan. Published Thesis.
8. Hessaen, M. H. & Fadlalmola, H. A. (2020). Assessment of Nursing Staff's Knowledge and Practice Regarding Streptokinase Administration for Myocardial Infarction Patients in Coronary Care Units in Public Hospitals. *European Journal of Pharmaceutical and Medical Research*, 7(7), 798-808.
9. Toppo, M., Dungdung, A. & Bakhla, A. (2019). Effectiveness of Teaching Program Regarding Thrombolytic Drug Therapy in Terms of Knowledge and Practice of Staff Nurses, *International Journal of Scientific Research*, 8 (4), 51-52 PRINT ISSN No 2277-8179.
10. Zakria, A. M., & Mohamed, S. A. (2017). Safety Intervention Educational Program to Reduce Medication Administration Errors and Interruptions. *IOSR Journal of Nursing and Health Science (IOSR-JNHS)*, 6(2), 15-25.
11. Slater, P., Edwards, R., & Badat, A., (2018). Evaluation of a staff well-being program in a pediatric oncology, hematology, and palliative care services group. *Journal of healthcare leadership*, 10, 67.